

Regained co-ordinated composure



CLINAZINE

(Trifluoperazine Hydrochloride B.P.)

For the treatment of anxiety in mild mental and emotional disturbances, and nausea and vomiting from a wide variety of causes.

CLINAZINE is rapid, effective, long acting, and exhibits a low incidence of side effects.

DOSE: General Practice—1 or 2 mg. b.i.d. Because of the long action of CLINAZINE, convenient twice daily dosage is generally sufficient and 4 mg. daily is usually adequate.

Psychiatric Practice—For the treatment of severe mental disorders requiring the higher dosage range, see CLINAZINE Medical Brochure available on request.

SIDE EFFECTS: General Practice—Daily doses of 2 to 4 mg. rarely produce serious side effects but drowsiness, dizziness, or stimulation may be observed. For description of possible neuromuscular (extrapyramidal) symptoms see CLINAZINE Medical Brochure available on request.

CONTRAINDICATIONS: CLINAZINE should not be administered to comatose or stuporous patients due to C.N.S. depressants. Use of the drug in cases of existing blood dyscrasias, bone marrow depression and pre-existing liver damage is contra-indicated.

SUPPLIED: In bottles of 50 sugar coated tablets 1 mg. (Yellow), 2 mg. (Green). Also available in tablets 5 mg. (Grey) and 10 mg. (Blue).

The CLINAZINE Medical Brochure is available on request for complete prescribing information.



MOWATT & MOORE LIMITED
Pointe Claire, Que., Canada